

Evaluation of Organic Soybean Varieties Southeast Research Farm, 2010

Kathleen Delate, professor
Jared Flater, research asst.
Departments of Horticulture and Agronomy
Kevin Van Dee, farm superintendent

Materials and Methods

According to the USDA National Organic Program, certified organic farmers must source organic seed (seed from organically raised crops). The organic seed industry is currently growing in Iowa and the Midwest. With this growth, organic growers are looking for University-based recommendations on organic varieties to use in Iowa. The Organic Agriculture Program at Iowa State University has been using organic seed at the Southeast Research Farm for eleven years with excellent results.

There were four varieties selected for the 2010 organic soybean variety trial. These included the following Blue River (Kelley, IA) 3.0 to 3.5–relative maturity hybrid soybean varieties: conventional, untreated BR35A0 was planted as a comparator against the three organic varieties, BR30A7, BR32F0, and BR34A7.

Plots measuring 20 x 185 ft. were laid out in a randomized complete block design with four replications of each variety. Winter rye was no-till drilled at a rate of 50 lbs/acre on April 2. Soybeans were planted at 160,000 seeds/acre at a depth of 1.5 in. on July 1, 2010. Weed management included rotary hoeing on July 14 and row cultivation on July 22, August 2 and August 19, 2010. Soybeans were harvested on October 22, 2010.

Plant stands and weed numbers were counted on July 26. Soybean cyst nematode samples were collected on September 21, and nematode analysis was conducted at the ISU Plant

Disease Clinic (Ames, IA). Grain samples were collected from each plot for grain quality analysis, which was conducted at the ISU Grain Quality Laboratory, Ames, IA.

Results and Discussion

Plant stands averaged 128,325 plants/acre in 2010, demonstrating a typical 20% reduction from planting rates after rotary hoeing and row cultivation tillage operations. Plant populations were greater in BR30A7 and 32F0 compared to BR34A7 and 35A0, averaging 136,700 plants/acre (Table 1). Weeds were well managed in 2010, with no significant differences among varieties. Grass weed populations averaged 6 weeds/sq. meter, and broadleaf weeds averaged 4 weeds/sq. meter (Table 1). Soybean cyst nematode populations averaged 671.88 eggs/100 cc of soil, with no significant differences among varieties. Organic soybean yields were excellent in 2010, averaging 42.13 bu/acre across all varieties, with a range of 37 to 45 bu/acre (Table 1). This yield was unexpectedly high, given the late planting date and the wet conditions in 2010.

Soybean grain quality was good considering the excess levels of moisture during the growing season, with no differences in moisture, protein and oil content among varieties (Table 2). Moisture content averaged 10.13% across all varieties (Table 2). Protein levels averaged 33.95% across all varieties (Table 2). Carbohydrate levels averaged 25.63%, with a lower level (24.4%) in BR30A7 (Table 2). Oil content averaged 17.5% across all varieties (Table 2). These results show great promise for organic seed. Because there was no statistical difference in yield and protein content between conventional and organic varieties, organic

producers can be confident that organic seed will produce at the same level as conventional seed.

Acknowledgments

We would like to thank the Leopold Center for Sustainable Agriculture for their support of this project. Thanks also go to Myron Rees and Chad Hesselteine, research farm staff, and Dan

Cwach, Jon Brunsvold, James Nguyen, and Vivian Bernau for their help in production, data collection and analytical aspects of this project. We also thank Charles Hurburgh and Glen Rippke of the ISU Grain Quality Lab, Kerry Culp of the ISU Soil and Plant Analysis Lab, and Blue River Hybrids for their support.

Table 1. Organic soybean stands, yield, soybean cyst nematode and weed populations at Southeast Research Farm, 2010.

Variety	Soybean stands (plants/acre)	Soybean yield (bu/ac)	Soybean cyst nematodes (eggs/100 cc soil)	Grass weeds (weeds/sq. meter)	Broadleaf weeds (weeds/sq. meter)
BR 30A7	137,400a	42.3	312.5	7.2	4.5
BR 32F0	136,000a	38.6	1262.5	4.5	4.0
BR 34A7	114,000c	43.0	500.0	6.8	3.7
35A0 Conv.	125,900b	44.6	612.5	6.5	3.1
LSD 0.05	9,100	NS	NS	NS	NS

Table 2. Soybean grain quality analysis–Southeast Research Farm, 2010.

Variety	Moisture (%)	Protein (%)	Oil (%)	Carbohydrates (%)
BR 30A7	9.6	35.3	17.4	24.4b
BR 32F0	9.5	34.8	17.1	25.1ab
BR 34A7	10.5	32.2	18.1	26.5a
35A0 Conv.	10.9	33.5	17.4	26.5a
LSD 0.05	NS	NS	NS	1.99