

Organic Corn Seed Coat Treatments–Neely-Kinyon Trial, 2006

Kathleen Delate, associate professor
Andrea McKern, research assistant
Departments of Horticulture and Agronomy
Bob Burcham, ag specialist

Introduction

Organic seed treatments have been limited in organic production, until recently. Agricoat Natural II™, is reported to be “an organic liquid biological seed treatment applied as a seed film coating and formulated with beneficial microbes, macro and micro nutrients, amino acids, organic acids, enzymes, proteins, vitamins, and minerals.” Natural II™ also is applied to “promote root growth and protect seedlings against early season soil-borne pathogens, improving germination, stand establishment and uniformity.”

Materials and Methods

Two treatments, replicated eight times in a completely randomized design, consisted of organic corn seed (BR 3448) treated with AgriCoat Natural II™ (Agricoat LLC, Soledad, CA) and a control (no seed treatment). Seed was planted on May 12, 2006, at 32,000 seeds/acre into a certified organic field that had received an application of hoop compost at 12 tons/acre on April 10, 2006. Plots were harrowed on May 22 and June 19, rotary hoed May 26 and June 1, and cultivated June 5, 2006. Plant stands were counted on June 8. Plots were harvested with a combine on November 1, 2006.

Results and Discussion

In 2006, there was no significant difference in corn stands between treatments (Table 1), despite an average increase of 333 plants per acre with the organic seed treatment. This contrasts with 2005 results, where stands were significantly greater in the treated plots. Yields were also not significantly different, with the coated seed plots yielding 194.5 bu/acre and the uncoated seed plots yielding 196.5 bu/acre (Table 1). These results may have been due to the warmer weather during the May planting of corn compared to cooler weather in 2005. Despite the lack of differences in 2006, yields from this organic variety were excellent, especially under the drought-like conditions of the 2006 season. Protein levels in harvested grain were also greater at 8.5% from the treated seed (Table 2).

Acknowledgments

We would like to thank the Leopold Center for Sustainable Agriculture. We also thank the Wallace Foundation and Practical Farmers of Iowa for their input and support. Thanks also go to David Rosmann, Magda Marin, Daniel Rosmann, and Greg Lilly; Todd Zehr of Agricoat LLC for providing the seed treatment, and Maury Johnson Blue River Hybrids for providing seed for this study.

Table 1. Corn stands and yields at Neely-Kinyon Farm, 2006.

Treatment	Corn stands plants/acre	Corn yield bu/acre
Coated seed	26,667	194.5
Uncoated seed	24,667	196.5
LSD (0.05)	NS	NS

Table 2. Corn grain quality at Neely-Kinyon Farm, 2006.

Treatment	Moisture %	Protein %	Oil %	Starch %
Coated seed	15.69	8.49a	3.61	59.97
Uncoated seed	15.86	8.31b	3.63	60.03
LSD (0.05)	NS	0.17	NS	NS