

Evaluation of Organic Soybean Rust Treatments for Organic Production —Neely-Kinyon Trial, 2007

Kathleen Delate, associate professor
Andrea McKern, research assistant
Departments of Horticulture and Agronomy
Bob Burcham, ag specialist

Introduction

Asian soybean rust, which arrived in the U.S. in 2004, has the potential to be the single most important impediment to economical organic production in the U.S., with the economic impact of ASR in organic systems ranging from \$30 to \$120 million in yield loss, upon its arrival in organic soybean areas. The fungus (*Phakopsora pachyrhizi*) survives year-round in warm areas, such as the southern United States. During a growing season, the pathogen is disseminated by northward seasonal wind.

Iowa State University has been awarded a grant from USDA to investigate “Strategies for Management of Asian Soybean Rust in Organic Systems,” which includes treatments allowable under certified organic conditions. Trials were established in 2005 in an area of Florida where rust is present to examine the effect of these treatments on ASR development and spread. Dry conditions across the south in 2007 kept soybean rust out of Iowa until very late in the season, when the first official find of ASR was reported in Adair County. Trials were established in Iowa, Pennsylvania and Michigan in 2006 and 2007 to examine yield effects of these treatments under non-rust conditions.

Materials and Methods

In the soybean rust treatment trial, Blue River 34A7 soybeans were planted at the Neely-Kinyon Farm on May 23, 2007, at 200,000 seeds/acre. Plots measuring 10 x 20 ft. were laid out in a randomized block design. There were four replications of the following treatments: MicroAF™ (TerraMax, Inc.,

Cottage Grove, MN) at 1 gallon/acre, Sonata® (AgraQuest, Inc., Davis, CA) at 1 gallon/acre, and Suregard Lime Sulphur Solution (Value Garden Supply, St. Joseph, MO) at 0.5 gallon/acre. All treatments were compared with a control. Treatments were applied on July 13, 2007, at the R-1 stage. Leaves were inspected on June 27 and July 12 for disease incidence before the treatments were sprayed, and on July 24, and August 6 and 20, by randomly selecting one leaf from the top, middle, and bottom sections of four plants per plot and recording incidence of disease. Plots were maintained with rotary hoeings on June 1, 8 and 19; cultivation on June 13 and 26, and July 26; and walking on July 17, 2007. Soybeans were harvested on October 27 with a combine. The percentage of stained soybeans in each treatment was determined by counting the number of stained soybeans in a 200-gram sample that was randomly collected from the harvest of each plot.

Results and Discussion

Soybean rust was not observed in the organic plots in 2007. Disease incidence was very low overall in 2007, ranging from 0 to 2.3% of leaves infected (Table 1). The treatments had no significant effect on yield, where soybeans averaged 63 bu/acre across all treatments (Table 1). There was no effect of the treatments on soybean diseases or grain quality (Table 1). Other diseases observed in 2007 include frog eye leaf spot, *Cercospora* leaf blight, bacterial pustule, bacterial blight, brown spot, and downy mildew. The most prevalent diseases were bacterial blight and bacterial pustule (Table 1 and Fig. 1). The highest number of infected leaves from bacterial blight occurred on July 12, while peak incidence of bacterial pustule was recorded on July 24. We will continue to test products in 2008, and monitor treatment effects

on all soybean diseases, in the event of rust appearing during the growing season in 2008.

References

<http://www.plantpath.iastate.edu/soybeanrust/>
<http://sbrusa.net>.
<http://extension.agron.iastate.edu/organicag/>

Acknowledgments

We would like to thank USDA-CSREES and the Leopold Center for Sustainable Agriculture

for their support of this project. Thanks also go to Greg Lilly, Mark Rosmann, Kelly Bevins, Chelsea Jensen, Ximena Cibils, Francisco Rosas and Francisco Viteri for their help. We also thank Blue River Hybrids, TerraMax, Inc., Cottage Grove, MN, AgraQuest, Inc., Davis, CA, and Value Garden Supply, St. Joseph, MO, and Charles Hurburgh and Glen Rippke of the ISU Grain Quality Lab for grain analysis.

Table 1. Yield and grain quality in soybean rust treatment trial, Neely-Kinyon, 2007.

Treatment	Yield (bu/acre)	Protein (%)	Oil (%)	Saturated Fatty Acids (%)	Carbohydrates (%)	Moisture (%)	Linolenic Acid (%)
Control	64.11 ^z	34.62	18.54	14.94	24.34	14.08	8.25
Sonata®	64.08	34.37	18.68	14.95	24.45	14.05	8.18
SureGard	64.98	34.19	18.71	14.78	24.60	13.97	8.18
MicroAF™	58.63	34.30	18.52	15.27	24.68	14.13	8.07
LSD 0.05	NS	NS	NS	NS	NS	NS	NS


^z Means within a column are not significant (NS), or significant at $p \geq 0.05$ (Fisher's protected LSD test).

Table 2. Disease incidence in soybean rust treatment trial, Neely-Kinyon, 2007.

Treatment	Average number of leaves infected (%)					
	Frogeye leaf spot	<i>Cercospora</i> leaf blight	Bacterial pustule	Bacteria I blight	Brown spot	Downy mildew
Control	0.00 ^z	0.80	1.50	2.30	0.80	0.80
Sonata®	1.50	0.00	1.50	2.30	0.00	1.50
SureGard	0.00	0.00	1.50	1.50	0.80	1.50
MicroAF™	0.15	0.00	0.00	1.50	0.00	0.00
LSD 0.05	NS	NS	NS	NS	NS	NS

^z Means within a column are not significant (NS), or significant at $p \geq 0.05$ (Fisher's protected LSD test).

Figure 1. Disease occurrence in soybean rust treatment trial, Neely-Kinyon, 2007.



* Data were taken before the treatment application on July 13.